

Excursion Set Theory for Correlated Random Walks

Arya Farahi¹ and Andrew J. Benson²

¹ *Michigan Center for Theoretical Physics, Randall Laboratory of Physics, The University of Michigan, Ann Arbor, MI 48109-1120, USA*

² *Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101, USA.*

5 March 2013

ABSTRACT

We present a new method to compute the first crossing distribution in excursion set theory for the case of correlated random walks. We use a combination of the path integral formalism of Maggiore & Riotto, and the integral equation solution of Zhang & Hui, and Benson et al. to find a numerically robust and convenient algorithm to derive the first crossing distribution in terms of a perturbative expansion around the limit of an uncorrelated random walk. We apply this methodology to the specific case of a Gaussian random density field filtered with a Gaussian smoothing function. By comparing our solutions to results from Monte Carlo calculations of the first crossing distribution we demonstrate that our method is accurate for power spectra $P(k) \propto k^n$ for $n = 1$, becoming less accurate for smaller values of n . It is therefore complementary to the method of Musso & Sheth, which will therefore be more useful for standard Λ CDM power spectra. Our approach is quite general, and can be adapted to other smoothing functions, and also to non-Gaussian density fields.

Key words: cosmology: theory, galaxies: haloes, dark matter, methods: analytical

1 INTRODUCTION

The growth of large scale structure in our Universe, and the subsequent formation of galaxies, is a challenging, non-linear problem. Small perturbations in the matter density in the early Universe grow via gravitational instability and eventually collapse to form dark matter halos. Haloes grow through merging with other halos in a hierarchy of ever increasing mass. This hierarchy of dark matter halos provides the cosmic web within which galaxy formation takes place. It is therefore crucial to understand the formation of dark matter halos in order to accurately model the formation of galaxies (Baugh 2006).

N-body simulations have developed into an extremely powerful tool to explore the formation and evolution of dark matter in the highly non-linear regime. However, they remain computationally expensive, making them impractical for surveying dark matter parameter space for example. An alternative approach to computing the formation of dark matter halos was first described by Press & Schechter (1974), and later developed extensively by Bond et al. (1991) and Lacey & Cole (1993), using excursion set theory. This extended Press-Schechter (ePS) methodology provides a highly successful statistical description of the formation histories of dark matter halos. For a recent review of excursion set theory see Zentner (2007).

The formation of halos has been studied using many different approaches, including numerical approaches such as Monte Carlo (e.g. Bond et al. (1991); Paranjape et al. (2012)), and analytical approaches (e.g. Bardeen et al. (1986); Bond et al. (1991); Maggiore & Riotto (2010a); Paranjape & Sheth (2012)). Of these analytic methods, the two most common are peaks theory, Bardeen et al. (1986), and the excursion set theory¹ Bond et al. (1991).

The excursion set theory used in the ePS technique follows random walks in the space of overdensity and variance (of the density field), and associates halo collapse with the first-crossing of a barrier by a random walk. For the case of a constant, or linear barrier and for random walks in which each step is uncorrelated with previous steps, an analytic solution can be found for the distribution of variances (and, therefore, masses) at which halo collapse occurs. This allows the halo mass function to be found directly for the simple case of a Gaussian-random density field and a variance derived using a filter that is sharp in k -space. Numerical methods exist to solve the problem numerically in the case of non-linear barriers (Zhang & Hui 2006;

¹ Recently, (Paranjape et al. 2012) and Paranjape & Sheth (2012) combined the excursion set and peak theory methods to construct an excursion set theory for peaks.

Benson et al. 2012), but non-Gaussian density fields and/or different filters, both of which cause successive steps in the random walk to become correlated, are more problematic.

Many attempts have been made to use path integral techniques—a powerful tool for studying stochastic and random processes—to solve problems in excursion set theory (e.g. Maggiore & Riotto (2010a,b,c); Ma et al. (2011); Adshead et al. (2012)). In this work, we modify the path integral method of Maggiore & Riotto (2010a) and combine it with our modified integral technique for solving the first crossing distribution, which is based upon the work of Zhang & Hui (2006) and which we introduced in our previous work, Benson et al. (2012). Through this combination, we develop a new approach to solve the first crossing problem in more general cases, and specifically address the case of a Gaussian filter.

The remainder of this work is organized as follows. In §2.2 we review how the path integral approach is used to find the probability distributions of trajectories and describe how to add a perturbation term representing the effects of the filter. In §3 we apply these methods to find the first crossing probability, while in §4 we demonstrate the method using a Gaussian filter. Finally in §5 we discuss our results.

We also include two appendixes. Appendix A presents a detailed calculation for the second-order contribution to the perturbation term and outlines how this same procedure can be used to derive higher order perturbation terms. Appendix B presents our numerical technique for solving the first crossing distribution which we have found to be robust against numerical error.

2 PROBABILITY DISTRIBUTION OF TRAJECTORY

2.1 Introduction

As shown by Maggiore & Riotto (2010a), Maggiore & Riotto (2010b), and Maggiore & Riotto (2010c) (hereafter MR10) in their series of works, path integral methods provide a powerful tool for studying the formation of dark matter halos in the context of excursion set theory. In their works they used this method to address the issues of non-Markovian trajectories, moving barriers, and non-Gaussianity in excursion set theory. Additionally, Zhang & Hui (2006) introduced an elegant method for solving the excursion set theory for moving barriers given knowledge of the probability distribution function for overdensities as a function of variance.

All of these methods have limitations, which we will discuss further in §5. In this work we modify the methods of MR10 and Zhang & Hui (2006) and combine them to solve the excursion set problem in the general case with greater accuracy. Specifically, our approach differs from that of MR10 as we find the probability distribution function for trajectories without consideration of the barrier, and then combine this with the method of Zhang & Hui (2006) to find the first crossing distribution for an arbitrary barrier.

In this section we review how the probability function for a trajectory can be found utilizing path integral techniques, and how one can find the relevant first-order perturbation term for different filters. In Appendix A we present a calculation of the second-order perturbation term, which, for many cases, will be sufficient.

2.2 Calculation of the Path Integral

We begin by applying the path integral method to the case of a Gaussian random density field with a sharp k -space filtering and demonstrate that in this case the well-known analytic solution is recovered.

We consider the density perturbation, $\delta(x)$, at point x in the universe, defined as

$$\delta(x) = \frac{\rho(x) - \bar{\rho}}{\bar{\rho}}. \quad (1)$$

In ePS theory we are interested in a smoothed density field, and so we smooth this density field with some filter function. Assuming a Gaussian-random density field, if the field is smoothed with a sharp k -space filter (i.e. a filter that is constant below some critical wavenumber, and zero otherwise) the resulting random walk in δ as a function of smoothing scale will be a Markovian process. This has a well-known analytic solution for the case of a linear barrier (Sheth 1998; Sheth & Tormen 2002).

After smoothing, the variance of density field is

$$S = \int d \ln k \Delta^2(k) |\tilde{W}^2(k, R)|, \quad (2)$$

where $\tilde{W}(k, R)$ is the filter function in Fourier space, R is the smoothing scale, and $\Delta(k)$ is the dimensionless power spectrum. In the path integral approach, we assume that $\delta(S)$ evolves stochastically with “time-like” S . By definition, $\langle \delta(S) \rangle = 0$, and for Gaussian fields the only non-vanishing connected correlator is the two-point correlator, $\langle \delta(S_1) \delta(S_2) \rangle$.

We wish to find the probability distribution of trajectories which start from $S = S_0$, $\delta = \delta_0$ and finish at $S = S_n$, $\delta = \delta_n$. We imagine discretizing our domain, $[S_0, S_n]$ into n equal sized parts. Each step is therefore $\epsilon = \Delta S/n$. The trajectory travels through S and in each step δ_i takes on a specific value. We will integral over all δ_i 's from $-\infty$ to ∞ to find the total probability for the trajectory given its starting and ending points. In the limit $n \rightarrow \infty$ we will recover the continuum solution. We define the probability density in the space of trajectories as

$$W(\delta_0; \delta_1, \dots, \delta_n; S_n) \equiv \langle \delta_D(\delta(S_0) - \delta_0) \cdots \delta_D(\delta(S_n) - \delta_n) \rangle, \quad (3)$$

where, for clarity, δ is the density fluctuation field and δ_D is the Dirac delta function. The probability distribution is found by integrating over all possible paths and has the form

$$P(\delta_0; S_0; S_1, \dots, S_{n-1}; \delta_n; S_n) = \int_{-\infty}^{+\infty} d\delta_1 \cdots \int_{-\infty}^{+\infty} d\delta_{n-1} \times W(\delta_0; \delta_1, \delta_2, \delta_3, \dots, \delta_{n-1}; \delta_n). \quad (4)$$

For the case of a Gaussian random density field and a sharp k -space filter this probability distribution depends only on the starting and ending points, as we will show below. The function $W(\delta_0; \delta_1, \dots, \delta_{n-1}; \delta_n)$ can be expressed in terms of the connected correlators of the theory,

$$W(\delta_0; \delta_1, \dots, \delta_{n-1}; \delta_n) = \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} \left[\frac{d\lambda_0}{2\pi} \cdots \frac{d\lambda_n}{2\pi} e^{i \sum_{i=1}^n \lambda_i \delta_i} \langle e^{-i \sum_{i=1}^n \lambda_i \delta(S_i)} \rangle \right], \quad (5)$$

or more practically,

$$W(\delta_0; \delta_1, \dots, \delta_{n-1}; \delta_n) = \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} \left[\frac{d\lambda_0}{2\pi} \cdots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i + \sum_{p=2}^{\infty} \frac{(-i)^p}{p!} \sum_{i_1=1}^n \cdots \sum_{i_p=1}^n \lambda_{i_1} \cdots \lambda_{i_p} \langle \delta_{i_1} \cdots \delta_{i_p} \rangle \right\} \right]. \quad (6)$$

Here $\langle \delta_{i_1} \cdots \delta_{i_p} \rangle$ denotes the connected n -point correlator. For a Gaussian random field density it is possible to show that the only non-vanishing connected correlator is the two-point correlator, $\langle \delta(S_1) \delta(S_2) \rangle$. Then for the Gaussian case we find,

$$W(\delta_0; \delta_1, \dots, \delta_{n-1}; \delta_n) = \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} \left[\frac{d\lambda_0}{2\pi} \cdots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i - \frac{1}{2} \sum_{i,j=1}^n \lambda_i \lambda_j \langle \delta_i \delta_j \rangle \right\} \right] \quad (7)$$

2.3 Gaussian Random Field with Sharp k -Space Filtering

It is well-known that for a Gaussian-random density field and sharp k -space filter the probability density function for trajectories is

$$P(\delta_0, S_0 = 0; \delta_n, S_n) = \frac{1}{\sqrt{2\pi S_n}} \exp \left\{ -\frac{(\delta_n - \delta_0)^2}{2S_n} \right\}. \quad (8)$$

Using eqns. (4) and (7) we should therefore be able to reproduce this known result. We find

$$P(\delta_0, S_0; \delta_n, S_n) = \int_{-\infty}^{+\infty} d\delta_1 \cdots \int_{-\infty}^{+\infty} d\delta_{n-1} \left[\int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} \frac{d\lambda_0}{2\pi} \cdots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i - \frac{1}{2} \sum_{i,j=1}^n \lambda_i \lambda_j \langle \delta_i \delta_j \rangle \right\} \right], \quad (9)$$

where $\langle \delta_i \delta_j \rangle$ is the two point correlation function. MR10 showed that for a Gaussian random field and sharp k -space filter the two-point correlator has the form,

$$\langle \delta(S_i) \delta(S_j) \rangle = \min(S_i, S_j) = \epsilon \min(i, j). \quad (10)$$

To solve the integrals in eqn. (9) we will use the following result (see, for example, Altland & Simons 2010),

$$\int d^n x \exp \left\{ -\frac{1}{2} \sum x_i A_{i,j} x_j + \sum b_i x_i \right\} = \frac{(2\pi)^{n/2}}{\sqrt{|A|}} \exp \left\{ \frac{1}{2} \sum b_i A_{ij}^{-1} b_j \right\}. \quad (11)$$

We first integrate over the λ 's in eqn. (9) by setting $b_i \equiv \delta_i$, $A_{ij} \equiv \langle \delta_i \delta_j \rangle$, and $x_i = \lambda_i$ to give²,

² For a detailed calculation, see MR10.

$$P(\delta_0; S_0 = 0; \delta_n; S_n) = \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \frac{1}{(2\pi\epsilon)^{n/2}} \exp \left\{ -\frac{1}{2\epsilon} \sum_{i=0}^{n-1} (\delta_{i+1} - \delta_i)^2 \right\}. \quad (12)$$

We now employ eqn. (9) again, this time with $x_i = \delta_i$. In this case, the elements of matrix A are $A_{i,i+1} = A_{i,i-1} = -1/\epsilon$ and $A_{i,i} = 2/\epsilon$, and the determinant of matrix A is $n\epsilon^{1-n}$. The elements of vector b are: $b_1 = b_{n-1} = -\epsilon^{-1}$ and the other elements are all zero. Using eqn. (11) to solve eqn (12) we find

$$P(\delta_0; S_0 = 0; \delta_n; S_n) = \frac{1}{\sqrt{2\pi n\epsilon}} \exp \left\{ -\frac{(\delta_n - \delta_0)^2}{2n\epsilon} \right\}. \quad (13)$$

Using the fact that $n\epsilon = S_n$ this reduces so,

$$P(\delta_0; S_0 = 0; \delta_n; S_n) = \frac{1}{\sqrt{2\pi S_n}} \exp \left\{ -\frac{(\delta_n - \delta_0)^2}{2S_n} \right\}, \quad (14)$$

which is independent of our discretization and agrees with the well-known analytic solution.

In the above we have assumed $S_0 = 0$, but in general, one may be interested to have the probability distribution of random walks starting from arbitrary S_0 . The two-point correlator is unchanged in this case, and so the above calculation holds, but now $n\epsilon = S_n - S_0$, and so

$$P(\delta_0; S_0; \delta_n; S_n) = \frac{1}{\sqrt{2\pi(S_n - S_0)}} \exp \left\{ -\frac{(\delta_n - \delta_0)^2}{2(S_n - S_0)} \right\}. \quad (15)$$

For solving the excursion set problem it will be useful to calculate the probability of a trajectory which starts at point $S_0 = 0$, $\delta = 0$ and then passes through the point $S = S_i$, $\delta = \delta_i$ before ending at $S = S_n$, $\delta = \delta_n$. We will use this result in section 3. In this case, we simply do not integrate over δ_i in eqn. (9),

$$P(\delta_0; \delta_i; S_i; \delta_n; S_n) = \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{i-1} \int_{-\infty}^{+\infty} d\delta_{i+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_i, \dots, \delta_n; S_n). \quad (16)$$

It is then straightforward to show that

$$P(\delta_0; \delta_i; S_i; \delta_n; S_n) = \frac{1}{\sqrt{4\pi^2 S_i (S_n - S_i)}} \exp \left\{ -\frac{(\delta_i - \delta_0)^2}{2S_i} \right\} \exp \left\{ -\frac{(\delta_n - \delta_i)^2}{2(S_n - S_i)} \right\}. \quad (17)$$

2.4 Gaussian Random Field with Arbitrary Filter

In this section we use the prescription introduced in §2.3 to calculate the probability function for trajectories with an arbitrary filter. For an arbitrary filter it is not possible to solve the necessary integrals analytically. Therefore, we adopt a perturbation theory approach to this problem. In this section we find the first-order solution, while Appendix A gives the second-order term. Finding higher order terms is possible, but in many cases second order will be sufficient.

Without loss of generality we will assume that $\delta_0 = 0$ and $S_0 = 0$. As we are still considering a Gaussian random field all correlations except the two-point correlation will vanish. We therefore assume that a small perturbation, $\Delta(S_i, S_j) \equiv \Delta_{ij}$, is added to the two-point correlation. We refer to Δ_{ij} as the “filter effect” term. We then have,

$$P(\delta_0; \delta_n; S_n) = \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \left[\int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{d\lambda_0}{2\pi} \dots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i - \frac{1}{2} \sum_{i,j=1}^n \lambda_i \lambda_j (\min(S_i, S_j) + \Delta_{ij}) \right\} \right]. \quad (18)$$

Calculation of Δ_{ij} for a given filter is discussed in § 2.5. Defining $\partial_i \equiv \partial/\partial\delta_i$, we can rewrite equation 18, retaining only first order corrections, as

$$P(\delta_0; \delta_n; S_n) = \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n) \quad (19)$$

$$+ \frac{1}{2} \sum_{i,j=1}^n \Delta_{ij} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \partial_i \partial_j W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n), \quad (20)$$

where,

$$W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n) = \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \left[\frac{d\lambda_0}{2\pi} \dots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i - \frac{1}{2} \sum_{i,j=1}^n \lambda_i \lambda_j (\min(S_i, S_j)) \right\} \right]. \quad (21)$$

Note that $\Delta_{ij} = \Delta_{ji}$ and $\Delta_{ii} = 0$. Since the two point correlation function is symmetric and we always have $\langle \delta_i \delta_i \rangle = \min(S_i, S_i) = S_i$, the perturbation to the two-point correlation function should be zero for $S_i = S_j$. Using $\Delta_{ii} = 0$ and the symmetry of the perturbation term one can write,

$$\frac{1}{2} \sum_{i,j=1}^n \Delta_{ij} \partial_i \partial_j = \frac{1}{2} \sum_{i=1}^{n-1} \Delta_{in} \partial_i \partial_n + \sum_{i=1}^{n-2} \sum_{j=i+1}^{n-1} \Delta_{ij} \partial_i \partial_j. \quad (22)$$

MR10 argued that $\sum_{i=1}^{n-1} \Delta_{in} \partial_i \partial_n$ gives an integral over an intermediate time, say S_j , and $\sum_{i=1}^{n-2} \sum_{j=i+1}^{n-1}$ produces a double integral over intermediate time, S . They therefore refer to these respectively as the “memory term” and “memory of memory term”. We will adopt the same labels in the remainder of this work.

Thus,

$$P(\delta_0; \delta_n; S_n) = P^{\text{gm}}(\delta_0; \delta_n; S_n) + P^{\text{mem}}(\delta_0; \delta_n; S_n) + P^{\text{mem-mem}}(\delta_0; \delta_n; S_n). \quad (23)$$

The probability distribution P^{gm} was derived in §2.3. We now consider the first perturbation term, P^{mem} :

$$P^{\text{mem}}(\delta_0; \delta_n; S_n) = \sum_{i=1}^{n-1} \Delta_{in} \partial_n \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \partial_i W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n). \quad (24)$$

In the continuum limit eqn. (24) this becomes an integral over time, S . We proceed by working with the discretized form and at the end will take the limit of $n \rightarrow \infty$ to find the continuum result:

$$\begin{aligned} P^{\text{mem}}(\delta_0; \delta_n; S_n) &= \sum_{i=1}^{n-1} \Delta_{in} \partial_n \lim_{\delta_c \rightarrow +\infty} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{i-1} \int_{-\infty}^{+\infty} d\delta_{i+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \\ &\quad \times W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_i = \delta_c, \dots, \delta_n; S_n) \\ &= \sum_{i=1}^{n-1} \Delta_{in} \partial_n \lim_{\delta_c \rightarrow +\infty} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{i-1} \int_{-\infty}^{+\infty} d\delta_{i+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \\ &\quad \times \frac{1}{(2\pi\epsilon)^{n/2}} \times \exp \left\{ -\frac{1}{2\epsilon} \sum_{j=0}^{i-1} (\delta_{j+1} - \delta_j)^2 \right\} \times \exp \left\{ -\frac{1}{2\epsilon} \sum_{j=i}^{n-1} (\delta_{j+1} - \delta_j)^2 \right\} \\ &= \sum_{i=1}^{n-1} \Delta_{in} \partial_n \lim_{\delta_c \rightarrow +\infty} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{i-1} \int_{-\infty}^{+\infty} d\delta_{i+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \\ &\quad \times \frac{1}{2\pi\epsilon} \times W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_i = \delta_c; S_i) \times W^{\text{gm}}(\delta_c; \delta_{i+1}, \dots, \delta_n; (S_n - S_i)), \end{aligned} \quad (25)$$

which can be simplified to give

$$P^{\text{mem}}(\delta_0; \delta_n; S_n) = \sum_{i=1}^{n-1} \Delta_{in} \partial_n \lim_{\delta_c \rightarrow +\infty} P^{\text{gm}}(\delta_0; \delta_i = \delta_c; S_i) P^{\text{gm}}(\delta_i = \delta_c; \delta_n; (S_n - S_i)). \quad (26)$$

It is obvious that for $i = 1, \dots, n-1$,

$$\partial_n \lim_{\delta_c \rightarrow +\infty} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{i-1} \int_{-\infty}^{+\infty} d\delta_{i+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \times W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_i = \delta_c, \dots, \delta_n; S_n) = 0. \quad (27)$$

Therefore, $P^{\text{mem}}(\delta_0; \delta_n; S_n) = 0$. In the same way, one can prove that $P^{\text{mem-mem}}(\delta_0; \delta_n; S_n) = 0$, implying that the whole perturbation term becomes zero as $\delta_c \rightarrow +\infty$. Consequently, it is possible to write the solution, when no barrier is considered, for all filters in terms of $\min(S_i, S_j) + \Delta_{ij}$, and all perturbation terms to the probability function become zero.

Now we again assume that the trajectory passes through point $S = S_k$, $\delta = \delta_k$. We will show that in this case the choice of filter does matter. For the memory term we find

$$P^{\text{mem}}(\delta_0; \delta_n; S_n) = \Delta_{kn} \partial_n \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{k-1} \int_{-\infty}^{+\infty} d\delta_{k+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \partial_k W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n), \quad (28)$$

where Δ_{kn} is the perturbation term and $\partial_k \equiv \partial/\partial\delta_k$ and $\delta_0 = 0$. This gives,




Figure 1. The FE for a Gaussian filter as given by eqn. (32). Blue, red, and green lines show the FE for $S_j = 1, 3, 5$ respectively.

$$\begin{aligned}
P^{\text{mem}}(\delta_0; \delta_n; S_n) &= \Delta_{kn} \partial_n \partial_k \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{k-1} \int_{-\infty}^{+\infty} d\delta_{k+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \frac{1}{(2\pi\epsilon)^{n/2}} \exp \left\{ -\frac{1}{2\epsilon} \sum_{i=0}^{n-1} (\delta_{i+1} - \delta_i)^2 \right\} \\
&= \Delta_{kn} \partial_n \partial_k \frac{1}{\sqrt{4\pi^2 S_k (S_n - S_k)}} \times \exp \left\{ -\frac{(\delta_k - \delta_0)^2}{2S_k} \right\} \times \exp \left\{ -\frac{(\delta_n - \delta_k)^2}{2(S_n - S_k)} \right\} \\
&= -\Delta_{kn} \frac{-\delta_k^2 S_n - S_k(\delta_n^2 - S_n + S_k) + \delta_k \delta_n (S_n + S_k)}{2\pi S_k^{3/2} (S_n - S_k)^{5/2}} \times \exp \left\{ -\frac{(\delta_k - \delta_0)^2}{2S_k} \right\} \times \exp \left\{ -\frac{(\delta_n - \delta_k)^2}{2(S_n - S_k)} \right\}.
\end{aligned} \tag{29}$$

For the memory-of-memory term we have,

$$P^{\text{mem-mem}}(\delta_0; \delta_n; S_n) = \Delta_{ij} \sum_{i < j} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{k-1} \int_{-\infty}^{+\infty} d\delta_{k+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \partial_i \partial_j W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n). \tag{30}$$

Therefore, all terms of $P^{\text{mem-mem}}$ vanish because there is at least one derivative with respect to δ_i for which $i \neq k$ which cancels the integral over δ_i . Taking the limit $\delta_i \rightarrow +\infty$ leaves us with zero.

2.5 Calculating the Filter Effect

We have not yet discussed how to calculate the filter effect (hereafter FE), $\Delta(S_i, S_j) = \Delta_{ij}$. A particular advantage of our method compared to its ancestors is that the terms that contribute to the FE are completely independent of the perturbation terms. This makes it very easy to develop solutions for alternative filters which can be used in eqn. (29).

The non-perturbative part of the two point correlation function was $\min(S_i, S_j)$. Therefore,

$$\Delta_{ij} = \langle \delta_i \delta_j \rangle - \min(S_i, S_j). \tag{31}$$

One can use the above formula to find the perturbation formula for any filter of interest. Note that, for most cases there will be no analytical solution, and the filter effect must be computed numerically for each point.

To be specific, we will now calculate the FE for a Gaussian filter. For simplicity, we consider a power spectrum of the form $P(k) \propto k^n$ with $n = 1$. Using eqn. (48) and assuming $i < j$ we find

$$\Delta_{ij}^{\text{GF}} = \frac{3S_i S_j - S_i^2 - 2\sqrt{(S_i^3 S_j)}}{S_i + S_j + 2\sqrt{S_i S_j}}, \tag{32}$$

where GF stands for ‘‘Gaussian filter’’. Note that Δ_{ij}^{GF} in eqn. (32) is not symmetric in its arguments as we have explicitly assumed that $i < j$.

In Fig. 1 we show the FE at three points. As expected, for small S the FE is less than $\min(S_i, S_j)$ and can be treated as a perturbation. For large variances the FE increases, so for large S the perturbation calculation will begin to break down.

3 FIRST CROSSING PROBABILITY

Zhang & Hui (2006) introduced an elegant method to find the first crossing probability in excursion set theory for arbitrary barriers. In our previous work (Benson et al. 2012) we modified this method to obtain more accurate answers for moving barriers. In this section we will incorporate the filter effect into this integral formulation. The method of Zhang & Hui (2006) begins by considering the integral form of the first crossing problem, which simply states that the total probability at any variance S is the sum of the probability that a trajectory has crossed the barrier at some smaller S , and the probability that the trajectory is currently below the barrier having never crossed at any smaller S :

$$1 = \int_0^{S_n} f(S') dS' + \int_{-\infty}^{B(S_n)} P_B(\delta, S_n) d\delta \quad (33)$$

where $P_B(\delta, S_n)$ is the distribution of δ at S_n accounting for the absorbing barrier and is given by

$$P_B(\delta_n, S_n) = P(\delta_n, S_n) - \int_0^{S_n} G(S_n, \delta_n, S') dS'. \quad (34)$$

In this formulation, eqns. (33) and (34), $G(S_n, \delta_n, S') dS'$ is the probability for a trajectory which crosses the barrier for the first time between S' and $S' + dS'$ and approaches the point (δ_n, S_n) . It is calculated by multiplying the fraction of trajectories which cross the barrier between S' and $S' + dS'$ for the first time, $f(S') dS'$, by the probability, $P(\delta_0; S_0; S', B(S'); \delta_n; S_n)$, of a trajectory which starts from point $(S_0 = 0, \delta = 0)$, crosses the barrier at point $(S = S', \delta = B(S'))$, and finally ends at point $(S = S_n, \delta = \delta_n)$. Here, $B(S)$, is the moving barrier. In general one can calculate the second term of equation 34 using,

$$G(S_n, \delta_n, S') \equiv f(S') \frac{P(\delta_0; S_0; S', B(S'); \delta_n; S_n)}{P(\delta_0; S_0; S', B(S'))}, \quad (35)$$

where $P(\delta_0; S_0; S', B(S'))$ is probability of trajectory which starts from point $(S = S_0, \delta = \delta_0)$ and ends at point $(S = S', \delta = B(S'))$. In the case of a Gaussian-random field and a sharp k -space filter one can divide eqn. (17) by eqn. (14) and obtain

$$G(S, \delta, S', B(S')) = f(S') \frac{1}{\sqrt{2\pi(S - S')}} \exp \left\{ -\frac{(\delta - B(S'))^2}{2(S - S')} \right\} = f(S') P_0(\delta - B(S'), S - S'). \quad (36)$$

For the case of a Gaussian-random field and a generic filter one can use eqn. (29) to obtain³,

$$\begin{aligned} G(S, \delta, S', B(S')) = & f(S') \frac{1}{\sqrt{2\pi(S - S')}} \exp \left\{ -\frac{(\delta - B(S'))^2}{2(S - S')} \right\} - f(S') \Delta(S, S') \\ & \times \frac{-B(S')S - S'(B(S) - S + S') + B(S')B(S)(S + S')}{\sqrt{2\pi S'^2(S - S')^5}} \exp \left\{ -\frac{(B(S) - B(S'))^2}{2(S - S')} \right\}. \end{aligned} \quad (37)$$

We define the second term in the above equation to be $f(S') \Delta(S', S) \gamma(S, \delta, S', B(S'))$. As we have shown in Benson et al. (2012), to discretize this equation and find the first crossing probability, $f(S)$, the next step is to find the following integral,

$$\Phi(S, S') \equiv \int_{-\infty}^{B(S)} \Delta(S', S) \gamma(S, \delta, S', B(S')) d\delta = -\Delta(S, S') \times \frac{B(S')S - B(S)S'}{\sqrt{2\pi S'^2(S - S')^3}} \exp \left\{ -\frac{(B(S) - B(S'))^2}{2(S - S')} \right\}. \quad (38)$$

We can now discretize eqn. (33) and solve for $f(S)$. In our previous work we showed in detail how one can discretize this equation. Briefly, we have

$$1 = \int_0^S f(S') dS' + \text{erf} \left\{ \frac{B(S)}{\sqrt{2S}} \right\} - \int_0^S f(S') \left[\text{erf} \left\{ \frac{B(S) - B(S')}{\sqrt{2(S - S')}} \right\} + 2\Phi(S, S') \right] dS'. \quad (39)$$

We assume that $\Delta S \equiv S_{\max}/N$. For eqn. (33) we find

³ Note that, in the general case, one should consider not the fraction of trajectories passing through point $(B[S], S)$ which end at point (δ_n, S_n) , but the fraction of trajectories which pass through point $(B[S], S)$ having never crossed the barrier for smaller S which end at point (δ_n, S_n) . For a Gaussian random field with a sharp k -space filter, these two are equivalent, as the path of any given trajectory for $S' > S$ is independent of its path for $S' < S$. For other filters, there is some correlation with the path for $S' < S$. Therefore, while eqn. (36) is exact, eqn. (37) is an approximation, obtained by ignoring the correlated nature of the random walk.

$$1 = \sum_{i=0}^{j-1} \frac{f(S_i) + f(S_{i+1})}{2} \Delta S + \operatorname{erf} \left\{ \frac{B(S_j)}{\sqrt{2(S_j)}} \right\} - \sum_{i=0}^{j-1} \left[f(S_i) \left(\operatorname{erf} \left\{ -\frac{B(S_j) - B(S_i)}{\sqrt{2(S_j - S_i)}} \right\} + 2\Phi(S_j, S_i) \right) + f(S_{i+1}) \left(\operatorname{erf} \left\{ \frac{B(S_j) - B(S_{i+1})}{\sqrt{2(S_j - S_{i+1})}} \right\} + 2\Phi(S_j, S_{i+1}) \right) \right] \frac{\Delta S}{2}. \quad (40)$$

Since $f(0) = 0$ we obtain,

$$\left[1 - \operatorname{erf} \left\{ \frac{B(S_j) - B(S_j)}{\sqrt{2(S_j - S_j)}} \right\} - 2\Phi(S_j, S_j) \right] \frac{\Delta S}{2} f(S_j) = 1 - \operatorname{erf} \left\{ \frac{B(S_j)}{\sqrt{2(S_j)}} \right\} - \sum_{i=1}^{j-1} f(S_i) \left(1 - \operatorname{erf} \left\{ \frac{B(S_j) - B(S_i)}{\sqrt{2(S_j - S_i)}} \right\} - 2\Phi(S_j, S_i) \right) \Delta S. \quad (41)$$

For all barriers of interest $\operatorname{erf} \left\{ \frac{B(S_j) - B(S_j)}{\sqrt{2(S_j - S_j)}} \right\} = 0$ and $\Phi(S_j, S_j) = 0$, such that

$$f(S_j) = \left[1 - \operatorname{erf} \left\{ \frac{B(S_j)}{\sqrt{2(S_j)}} \right\} - \sum_{i=1}^{j-1} f(S_i) \left(1 - \operatorname{erf} \left\{ \frac{B(S_j) - B(S_i)}{\sqrt{2(S_j - S_i)}} \right\} - 2\Phi(S_j, S_i) \right) \Delta S \right] \frac{2}{\Delta S}. \quad (42)$$

In the above, $\Phi(S_j, S_i)$ plays the role of a filter correction term, and is calculated using eqn. (38).

Here we have computed up to first-order perturbations. In Appendix A we derive results up to second-order which is sufficient for most purposes (although this will depend on the choice of filter). We address numerical issues in solving for $f(S_j)$ in Appendix B.

3.1 Gaussian Random Field with Gaussian filter

So far, our calculations are independent of the choice of filter. In this section we compute the first crossing distribution for a Gaussian filter applied to a Gaussian-random field. For a Gaussian-random field both the unconditional and conditional probabilities should also be Gaussian, independent of the choice of filter. We use this case to illustrate that the path integral formalism gives the expected answer. Of course, the true power of the path integral formalism will become relevant when applied to non-Gaussian fields. The Gaussian filter is convenient as its two point correlation function can be calculated analytically and, unlike the sharp k -space filter, its smoothing region in configuration space is well defined and localized. In this case, the variance, S , is defined as

$$S = \langle \delta^2(R) \rangle = \frac{1}{2\pi^2} \int k^2 P(k) \tilde{W}^2(k, R) dk, \quad (43)$$

where

$$\tilde{W}(k, R) = \exp \left\{ -\frac{R^2 k^2}{2} \right\} \quad (44)$$

is the Fourier transform of our filter which in real space has the form

$$W(r, R) = \frac{1}{(2\pi R^2)^{3/2}} \exp \left\{ -\frac{r^2}{2R^2} \right\}. \quad (45)$$

For a scale-free power spectrum, $P(k) = Ak^n$,

$$S = \langle \delta^2(R) \rangle = \frac{A}{4\pi^2} \Gamma \left(\frac{3+n}{2} \right) R^{-3-n}, \quad (46)$$

and two-point correlation function is

$$\langle \delta(R_1) \delta(R_2) \rangle = \frac{A}{4\pi^2} \Gamma \left(\frac{3+n}{2} \right) \left(\frac{R_1^2}{2} + \frac{R_2^2}{2} \right)^{-\frac{3-n}{2}}, \quad (47)$$

where here Γ is the usual Gamma function. For $n = 1$ therefore, we find the following relation between the variance and the two-point correlation function:

$$\langle \delta_i \delta_j \rangle = \frac{4S_i S_j}{S_i + S_j + 2\sqrt{S_i S_j}}. \quad (48)$$

For general exponent n this becomes:

$$\langle \delta_i \delta_j \rangle = \left(\frac{S_i^{-\frac{2}{3+n}} + S_j^{-\frac{2}{3+n}}}{2} \right)^{-\frac{3+n}{2}}. \quad (49)$$

Setting $i = j$ gives us the variance of point S_i , $\langle \delta_i \delta_j \rangle = S_i$, consistent with that found for a sharp k -space filter for which $\langle \delta_i \delta_j \rangle = \min(S_i, S_j)$.

4 RESULTS

We can now use our results to compute the first crossing distribution for a Gaussian filter. MR10 also employed the path integral method account for the filter effect as perturbation term. They used an approximate form for the two-point correlation function for both a top-hat filter in real space and a Gaussian filter. Specifically, they proposed the following function:

$$f(S) = \frac{1 - \kappa}{\sqrt{2\pi}} \frac{\delta_c}{S^{3/2}} \exp \left\{ -\frac{\delta_c^2}{2S} \right\} + \frac{\kappa}{2\sqrt{2\pi}} \frac{\delta_c}{S^{3/2}} \Gamma \left(0, \frac{\delta_c^2}{2S} \right), \quad (50)$$

where $\Gamma(0, z)$ is the incomplete Gamma function, $\delta_c \approx 1.686$ is the barrier for spherical collapse. MR10 computed the value of κ for top-hat and Gaussian filters, and a Λ CDM power spectrum, finding that $\kappa \approx 0.4592$ and 0.35 respectively with a weak dependence on scale. Because MR10 computed κ for a Λ CDM power spectrum it is not directly comparable to the power-law cases we consider here. Furthermore, for case of power-law, we have compared the form of our expression for Δ_{ij} and find that it is not well approximated by MR10's fitting function for any value of κ . Therefore, we instead compare our solutions to the results of Monte Carlo simulations of the first crossing problem.

Peacock & Heavens (1990) used an alternative method to find the first crossing probability distribution for correlated random walks, proposing an approximate formula for the Gaussian filter. Later Paranjape & Sheth (2012) proposed another approximate formula and argued that their formula worked well for a large range of S . More recently, MS12 used this method to propose an improved approximation, which is similar to that of Peacock & Heavens (1990), for generic filters as well as moving barriers.

Fig. 2 compares results from this work with those of MS12, and a Monte Carlo calculation of trajectories⁴. We use Robertson et al. (2009)'s prescription (see also Bond et al. 1991) for Monte Carlo simulation of random walks. Results are shown for two different values of the power spectrum index, n . For $n = +1.0$ (left panel) our results are in very good agreement with the Monte Carlo results for $\nu \gtrsim 1$ when we include up to the second-order perturbation term. For smaller ν , our result begins to diverge from the Monte Carlo result, albeit slowly. Higher order perturbation terms could be included of course, which would extend the validity of our result to smaller ν . The approximation of MS12 performs poorly for $n = +1.0$, differing significantly from the Monte Carlo solution for $\nu \lesssim 6$. For $n = -1.2$ the situation is reversed. Here, the approximation of MS12 works extremely well (as MS12 showed), while our approximation disagrees substantially with the Monte Carlo result.

Our method applies equally well to non-constant (a.k.a. “moving”) barriers. Figure 3 compares results from this work with those of MS12, and a Monte Carlo calculation of trajectories for a linear moving barrier with the form of,

$$B(S) = \delta_c \left(1 + \frac{\alpha S}{\delta_c^2} \right). \quad (51)$$

The figure shows the result for two choices of α , $+0.5$ (dashed lines) and -1.0 (solid lines), and two different power spectrum, $n = +1.0$ (left panel) and $n = -1.2$ (right panel). For both power spectra, the approximation of MR12 and this are in close agreement for $\alpha < 0$ (although MS12 remains the better match to the Monte Carlo results for $n = -1.2$), but as α is increased they diverge from each other, such that for $n = +1.0$ (left panel) our results are in very good agreement with the Monte Carlo results, while for $n = -1.2$ the approximation of MS12 works better (as MS12 showed).

⁴ We cannot compare with the results of MR10 in this case. Their mass function is expressed in terms of a parameter, κ , which parameterizes the perturbation, Δ_{ij} . We find that the parameterization used by MR10—derived for a Λ CDM power spectrum—is not a good match to the perturbation derived in this work for any value of κ . Note that MR10 used transfer function for calculating κ but in this work just the power-law is considered. For similar reasons, the work of Corasaniti & Achitouv (2011) and Achitouv & Corasaniti (2012) are also not comparable to our current examples.




Figure 2. The dimensionless halo mass function $\nu f(\nu)$ as computed using different approximations, for a Gaussian-random density field, smoothed with a Gaussian filter, and for two different power spectral indices: $n = +1.0$ (left panel) and $n = -1.2$ (right panel). The dark blue line shows the results from a Monte Carlo calculation, while red and magenta lines show results from this work including up to the first and second-order perturbations respectively. Black lines show the approximation of MS12. For reference, the green line shows the exact result for a sharp k -space filter.




Figure 3. The dimensionless halo mass function $\nu f(\nu)$ as computed using different approximations, for a Gaussian-random density field, smoothed with a Gaussian filter, and for two different power spectral indices: $n = +1.0$ (left panel) and $n = -1.2$ (right panel), for the case of a linear moving barrier of the form $B(S) = \delta_c(1 + \alpha S/\delta_c^2)$. Dashed lines show results for $\alpha = +0.5$ while solid lines show results for $\alpha = -1.0$. The dark blue line shows the results from a Monte Carlo calculation, while red and magenta lines show results from this work including up to the first and second-order perturbations respectively. Black lines show the approximation of MS12. For reference, the green line shows the exact result for a sharp k -space filter.

5 CONCLUSION AND DISCUSSION

Many attempts have been made to account for correlated random walks in excursion set theory, whether induced by the filter effect or by intrinsic correlations in the density field. For example, Peacock & Heavens (1990) computed approximate solutions for correlated random walks, specifically for a Gaussian filter function. This approach was later improved (Paranjape & Sheth 2012; Paranjape et al. 2012; Musso & Sheth 2012; Musso et al. 2012) to work with moving barriers. While direct Monte Carlo simulation can account for these correlations it is inefficient and therefore impractical for application within semi-analytical models which use ePS theory to construct merging histories for dark matter halos. MR10 employed a path integral method to include the filter effect as a perturbation term. They used an approximate function for the two-point correlation function arising from both real-space top-hat, and Gaussian filters. In their later works they have proposed a series solution for moving barriers with sharp k -space filters. It is difficult to include both the filter effect and a moving barrier in the approach of MR10.

Later works (Corasaniti & Achitouv 2011) improved MR10’s method by considering a linear barrier in the perturbative path integral method, but still are not applicable to an arbitrary moving barrier.

The solution for arbitrary moving barriers with uncorrelated random walks is well understood and has been studied for many years (Zentner 2007). In particular, the integral method of Zhang & Hui (2006), and its numerically more robust modification (Benson et al. 2012), work well in this case. In this work we have combined the power of the path-integral approach to excursion set theory with the integral-method solution for arbitrary barriers to develop a perturbative solution to the first crossing problem applicable to arbitrary filters and barriers. The perturbation terms can be computed directly given the functional form of the filter. Our approach utilizes the path integral formalism of MR10, but uses it to find the unconditional probability $P_0(\delta, S)$ —the probability that a random walk reaches the point δ at variance S —and the conditional probability $P(\delta_c, S_c; \delta, S)$ —the probability that a random walk cross the barrier at point S then reaches the point δ at variance S —considering all possible walks and weighting them equally when constructing the ensemble average. Note that we specifically *do not* use the path integral formalism to compute the probability $P(\delta, S)$ in the presence of the absorbing barrier. In this formalism, the conditional probability is responsible for accounting for correlations between steps in the random walk. Given these unconditional and conditional probabilities, the barrier is added to the problem and used to compute the first crossing distribution using the method of Zhang & Hui (2006). We find that this approach is accurate when compared to the results of Monte Carlo simulations over a wide range of variance.

Our method complements that of MS12, which is designed to be accurate in the limit of smooth random walks (corresponding to more negative values of the spectral index n). Our approach expands around the limit of uncorrelated steps in the random walk (corresponding to “jagged” random walks, and more positive values of the spectral index n). This can be clearly seen in Figure 3 in which we show results for $n = -1.2$ and $n = +1.0$. The MS12 approach is therefore more useful for standard Λ CDM power spectra (for which the local exponent is negative on scales of interest for dark matter halo formation). Finally, we compared our solution with that of MS12 for a linear moving barrier with the form $B(S) = \delta_c(1 + \alpha S/\delta_c^2)$ for two choice of α , $\alpha = +0.5$ and $\alpha = -1.0$. As can be seen in Figure 2, for $\alpha < 0$ both methods are in approximate agreement with Monte Carlo calculations.

Our method has the advantage of being very flexible. Currently, we have used these methods to derive the unconditional and conditional probability distributions (required to solve the first crossing problem) for the simple case of a Gaussian random field—demonstrating that this approach correctly recovers the expected Gaussian distributions in this case. In a future work, we intend to exploit this flexibility of our method to examine models in which the density field is non-Gaussian. Additionally, we note that our approach currently gives equal weight to each possible random walk when constructing the ensemble average. Dark matter halos may, in fact, correspond to a special subset of all walks, e.g. those corresponding to peaks in the density field (Paranjape & Sheth 2012; Paranjape et al. 2012), which would require different weights to be assigned to random walks in the ensemble averaging. We plan to revisit this issue.

ACKNOWLEDGMENTS

We thank Ravi Sheth, Brant Robertson, and Antonio Riotto for helpful discussions.

REFERENCES

- Achitouv I., Corasaniti P., 2012, JCAP, 1202, 002
- Adsheed P., Baxter E. J., Dodelson S., Lidz A., 2012, Phys.Rev., D86, 063526
- Altland A., Simons B., 2010, Condensed Matter Field Theory. Cambridge University Press
- Bardeen J. M., Bond J. R., Kaiser N., Szalay A. S., 1986, ApJ, 304, 15
- Baugh C. M., 2006, Rept.Prog.Phys., 69, 3101
- Benson A. J., Farahi A., Cole S., Moustakas L. A., Jenkins A., et al., 2012, MNRAS
- Bond J. R., Cole S., Efstathiou G., Kaiser N., 1991, ApJ, 379, 440
- Corasaniti P., Achitouv I., 2011, Phys.Rev., D84, 023009
- Lacey C., Cole S., 1993, Monthly Notices of the Royal Astronomical Society, 262, 627
- Ma C.-P., Maggione M., Riotto A., Zhang J., 2011, MNRAS, 411, 2644
- Maggione M., Riotto A., 2010a, ApJ, 711, 907
- , 2010b, ApJ, 717, 515
- , 2010c, ApJ, 717, 526
- Musso M., Paranjape A., Sheth R. K., 2012, MNRAS
- Musso M., Sheth R. K., 2012, MNRAS
- Paranjape A., Lam T. Y., Sheth R. K., 2012, MNRAS, 420, 1648
- Paranjape A., Lam T. Y., Sheth R. K., 2012, MNRAS, 420, 1429

- Paranjape A., Sheth R. K., 2012, MNRAS, 419, 132
 Paranjape A., Sheth R. K., 2012, MNRAS, 426, 2789
 Paranjape A., Sheth R. K., Desjacques V., 2012, ArXiv e-prints
 Peacock J., Heavens A., 1990, MNRAS, 243, 133
 Press W. H., Schechter P., 1974, ApJ, 187, 425
 Robertson B. E., Kravtsov A. V., Tinker J., Zentner A. R., 2009, ApJ, 696, 636
 Sheth R. K., 1998, MNRAS, 300, 1057
 Sheth R. K., Tormen G., 2002, MNRAS, 329, 61
 Zentner A. R., 2007, Int.J.Mod.Phys., D16, 763
 Zhang J., Hui L., 2006, ApJ, 641, 641

APPENDIX A: CALCULATION OF THE PERTURBATION

In § 2.4 we described in detail the calculation of the first-order perturbation arising from the filter effect. Below we outline the calculation of the second-order perturbation term. The same approach can be used to derive yet higher order perturbations if necessary. We have found that, for large masses, the first perturbation term is sufficient. Adding the second or higher order terms extends the range of validity of our approximations to larger S (smaller mass).

Again we assume that $\delta_0 = 0$ and $S_0 = 0$, and assume a density field with Gaussian-random fluctuations. Therefore, the probability of finding the trajectory at (δ_n, S_n) is,

$$P(\delta_0; \delta_n; S_n) = \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \left[\int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{d\lambda_0}{2\pi} \dots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i - \frac{1}{2} \sum_{i,j=1}^n \lambda_i \lambda_j (\min(S_i, S_j) + \Delta_{ij}) \right\} \right], \quad (\text{A1})$$

where Δ_{ij} is the filter effect(FE) term. By defining $\partial_i \equiv \partial/\partial\delta_i$, we can write equation A1, including perturbations up to second-order as,

$$\begin{aligned} P(\delta_0; \delta_n, S_n) &= \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n) \\ &+ \frac{1}{2} \sum_{i,j=1}^n \Delta_{ij} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \partial_i \partial_j W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n) \\ &+ \frac{1}{2!} \times \frac{1}{4} \sum_{i,j,k,m=1}^n \Delta_{km} \Delta_{ij} \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \partial_i \partial_j \partial_k \partial_m W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n) \\ &+ \dots, \end{aligned} \quad (\text{A2})$$

where,

$$W^{\text{gm}}(\delta_0; \delta_1, \dots, \delta_n; S_n) = \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \left[\frac{d\lambda_0}{2\pi} \dots \frac{d\lambda_n}{2\pi} \exp \left\{ i \sum_{i=1}^n \lambda_i \delta_i - \frac{1}{2} \sum_{i,j=1}^n \lambda_i \lambda_j (\min(S_i, S_j)) \right\} \right]. \quad (\text{A3})$$

Note again that the perturbation term has the following properties: $\Delta_{ij} = \Delta_{ji}$, and $\Delta_{ii} = 0$. Because the two point correlation function is symmetric and $\langle \delta_i \delta_i \rangle = \min(S_i, S_i) = S_i$, the perturbation part of two point correlation function is always zero. Using $\Delta_{ii} = 0$ and the symmetry of perturbation term one can write,

$$\frac{1}{2} \sum_{i,j=1}^n \Delta_{ij} \partial_i \partial_j = \frac{1}{2} \sum_{i=1}^{n-1} \Delta_{in} \partial_i \partial_n + \sum_{i=1}^{n-2} \sum_{j=i+1}^{n-1} \Delta_{ij} \partial_i \partial_j, \quad (\text{A4})$$

and,

$$\frac{1}{4} \sum_{i,j,k,m=1}^n \Delta_{ij} \Delta_{km} \partial_i \partial_j \partial_k \partial_m = \left(\frac{1}{2} \sum_{i=1}^{n-1} \Delta_{in} \partial_i \partial_n + \sum_{i=1}^{n-2} \sum_{j=i+1}^{n-1} \Delta_{ij} \partial_i \partial_j \right) \times \left(\frac{1}{2} \sum_{k=1}^{n-1} \Delta_{kn} \partial_k \partial_n + \sum_{k=1}^{n-2} \sum_{m=k+1}^{n-1} \Delta_{km} \partial_k \partial_m \right). \quad (\text{A5})$$

In § 2.4 we demonstrated that after integration all terms vanish when finding the probability of a trajectory that ends at (δ_n, S_n) , but when considering a trajectory that passes through (δ_k, S_k) as an intermediate step the term with $\Delta_{kn} \partial_n \partial_k$ is non-vanishing for the first-order perturbation. We apply this same argument for the second-order perturbation term and find

that the only non-vanishing term is that with $\Delta_{kn}^2 \partial_n^2 \partial_k^2$. For higher-order perturbations the same argument can be used to show that the only non-vanishing term in the N^{th} -order perturbation is: $\Delta_{kn}^N \partial_n^N \partial_k^N$.

We define the N^{th} -order perturbation term for the trajectory that passes (δ_k, S_k) and ends at (δ_n, S_n) as:

$$\begin{aligned} P_{\text{per}}^N(\delta_0; \delta_k; S_k; \delta_n; S_n) &= \frac{1}{N!} \Delta_{kn}^N \partial_n^N \partial_k^N \int_{-\infty}^{+\infty} d\delta_1 \dots \int_{-\infty}^{+\infty} d\delta_{k-1} \int_{-\infty}^{+\infty} d\delta_{k+1} \dots \int_{-\infty}^{+\infty} d\delta_{n-1} \\ &\quad \times \frac{1}{(2\pi\epsilon)^{n/2}} \exp \left\{ -\frac{1}{2\epsilon} \sum_{i=0}^{n-1} (\delta_{i+1} - \delta_i)^2 \right\} \\ &= \frac{1}{N!} \Delta_{kn}^N \partial_n^N \partial_k^N \frac{1}{\sqrt{4\pi^2 S_k (S_n - S_k)}} \times \exp \left\{ -\frac{(\delta_k - \delta_0)^2}{2S_k} \right\} \times \exp \left\{ -\frac{(\delta_n - \delta_k)^2}{2(S_n - S_k)} \right\}. \end{aligned} \quad (\text{A6})$$

The factor $1/N!$ in P_{per}^N arises from the Taylor expansion of eqn.(A1).

To find $\Phi(S, S')$, as defined in eqn. (38) we employ eqn. (36) to integrate with respect to δ from $-\infty$ to $B(S)$, and set $\delta_k = B(S')$ and $S_k = S'$. For the N^{th} -order perturbation we find:

$$\Phi^N(S, S') \equiv \int_{-\infty}^{B(S)} \sqrt{2\pi S'} \exp \left(\frac{\delta'}{2S'} \right) P_{\text{per}}^N(\delta_0; \delta_k; S_k; \delta_n; S_n) d\delta. \quad (\text{A7})$$

Specifically, the first and second-order perturbation terms are therefore found to be,

$$\Phi^1(S, S') = -\Delta(S, S') \times \exp \left\{ -\frac{(B(S) - B(S'))^2}{2(S - S')} \right\} \times \frac{B(S')S - B(S)S'}{\sqrt{2\pi S'^2 (S - S')^3}}, \quad (\text{A8})$$

and,

$$\begin{aligned} \Phi^2(S, S') &= \frac{\Delta(S, S')}{2} \times \exp \left\{ -\frac{(B(S) - B(S'))^2}{2(S - S')} \right\} \\ &\quad \times \left[\frac{-B^3(S)S'^2 + B^2(S)B(S')S'(S' + 2S) - B(S)(2S' + S)(S'^2 + B^2(S')S - S'S) + B(S')S(3S'(S' - S) + B^2(S')S)}{\sqrt{2\pi S'^4 (S - S')^7}} \right], \end{aligned} \quad (\text{A9})$$

respectively.

APPENDIX B: NUMERICAL ALGORITHM

In our previous work (Benson et al. 2012) and here in §3 we employ a first order finite difference method to discretize and solve the integral equation (39). The solution to this equation contains high-order cancellations which can make it numerically challenging to solve. We have found that for a sharp k -space filter this method is robust for all S and for arbitrary barriers providing that care is taken in the numerical evaluation. However, we find that the addition of perturbation terms can induce numerical inaccuracies that lead to fluctuation and divergence of the solution for $S > 1$. To circumvent this problem we proceed as follows.

We begin by rewriting eqn. (41) in a more general form, considering all perturbation terms:

$$\begin{aligned} 1 - \text{erf} \left\{ \frac{B(S_j)}{\sqrt{2(S_j)}} \right\} &= \left[1 - \text{erf} \left\{ \frac{B(S_j) - B(S_j)}{\sqrt{2(S_j - S_j)}} \right\} - 2 \sum_{k=1}^N \Phi^k(S_j, S_j) \right] \frac{\Delta S}{2} f(S_j) \\ &\quad + \sum_{i=1}^{j-1} f(S_i) \left(1 - \text{erf} \left\{ \frac{B(S_j) - B(S_i)}{\sqrt{2(S_j - S_i)}} \right\} - 2 \sum_{k=1}^N \Phi^k(S_j, S_i) \right) \Delta S. \end{aligned} \quad (\text{B1})$$

We now adjust the weight given to the first term on the right-hand side of eqn. (41) from $\Delta S/2$ to $\Delta S/\alpha$. Solving for $f(S_j)$ gives

$$f(S_j) = \left[1 - \text{erf} \left\{ \frac{B(S_j)}{\sqrt{2(S_j)}} \right\} - \sum_{i=1}^{j-1} f(S_i) \left(1 - \text{erf} \left\{ \frac{B(S_j) - B(S_i)}{\sqrt{2(S_j - S_i)}} \right\} - 2 \sum_{k=1}^N \Phi^k(S_j, S_i) \right) \Delta S \right] \frac{\alpha}{\Delta S}. \quad (\text{B2})$$

Note that $f(S_j)$ is proportional to α . However, all $f(S_{i < j})$ also depend on α . In particular, if we make α slightly smaller than 2 we cause successive $f(S_j)$ to oscillate around the true solution, effectively damping the numerical noise which otherwise causes divergence.

Figures B1 and B2 illustrate the convergence achieved using this numerical recipe. In both cases we compare our numerical




Figure B1. The relative error of our numerical method with respect to the analytic solution for a sharp k -space filter, Gaussian-random density field, and constant barrier. Results are shown for different values of the numerical parameter, α . The mesh size is fixed at 600 in all cases.

results with the analytic solution for a sharp k -space filter, a Gaussian random density field and a constant barrier. Figure B1 shows the relative error in the numerical and analytic solution for different values of α and a fixed mesh size of 600. While the error is lowest for $\alpha = 2$ over most of the range, it is clearly growing for the largest values of S . We find that it continues to do so as S is increased further. As our solution is typically valid only for small S this may not be a problem. Nevertheless, we find that adopting a value of α slightly less than 2 results in acceptably small errors, and an error term that decreases with increasing S . Figure B2 considers different mesh sizes for fixed $\alpha = 1.5$. The numerical solution is stable (with increasing accuracy) as the mesh size is increased, demonstrating that our solutions are robust and numerically converged.




Figure B2. The relative error of our numerical method with respect to the analytic solution for a sharp k -space filter, Gaussian-random density field, and constant barrier. Results are shown for different mesh sizes (as indicated in the figure; here n is the number of mesh points in the range $S = [0, 10]$), and for a fixed $\alpha = 1.5$.